## Items and Scoring for the Substance Use Risk Profile-Pregnancy Scale State at the beginning that the patient's

responses are confidential (Refer to script)

1) Low risk if they never used alcohol; 2) average risk if they had used alcohol in the past but consumed no alcohol and smoked fewer than three cigarettes in month before pregnancy: 3) high risk if they consumed any alcohol or smoked at least three

cigarettes in the month before pregnancy
Items:  1. Have you ever smoked marijuana?  No Yes
2. In the month before you knew you were pregnant, how many beers, how much wine, or how much liquor did you drink?  None  Yes,
<ul> <li>Have you ever believed that you needed to cut down on your alcohol and or drug use (including the non-medical use of prescription medication)?</li> <li>No Yes</li> </ul> Scoring:
Classify the number of alcoholic drinks before pregnancy as none compared to any, and count the number of affirmative items.
0 = low risk 1 = moderate risk 2-3 = high risk
In low-risk populations, one or more affirmative items indicate a positive screen, whereas, in high-risk populations, two or more affirmative items ndicate a positive screen.
Staff Signature: Date:
Interpreter Used: □ No □ Yes Interpreter's Name:
275 East Main St. Phone: (859) 625-8466

\*Screening for prenatal substance use: Development of the Substance Use Risk Profile-Pregnancy scale. Yonkers et al. Obstet Gynecol 2010; 116:827-33

Kentucky Perinatal Quality Collaborative

**Email: KyPQC@ky.gov** 

Frankfort, Kentucky 40621

## Substance Use Screener Follow-Up Questions

Women who answered yes to any of the 5Ps should be further assessed for opioid or any other drug use.

1. Have you used opioids, narcotics, or pain medication in the last year?
No Yes
2. If yes, please define the length of time and frequency:
Length of time and frequency:
3. If yes, were they prescribed or unprescribed?
Prescribed Unprescribed
4. Have you used any other drugs or unprescribed medications in the last year?
No Yes
Patients with positive responses who have used unprescribed drugs in pregnancy or used prescribed opioids for longer than a month need the referral protocol to link them to services and medication for opioid use disorder (MOUD), if indicated.
(For positive responses to 5Ps, also document that Kentucky All Schedule Prescription Electronic Report [KASPER] has been reviewed.)

275 East Main St. Frankfort, Kentucky 40621 Phone: (859) 625-8466 Email: KyPQC@ky.gov

